


Feed the World in the Face of Numerous Destabilizing Factors

- Changing climates
- Competing water demands
- Pest resistance to chemicals
- Declining soil and water quality











Climate Smart Farming Precision Agriculture (site specific and adaptive)



Translational Pipelines

Realm of science

Realm of practice




Precision Sustainable Agriculture and GROW Networks:


Transdisciplinary Research & Translational Pipelines

Precision Sustainable Agriculture


Precision Sus




On-farm and research stations LIVING LABORATORIES SOIL PLANT SENSORS PHENO'STRESS' CAMS PLANT SENSORS PHENO'STRESS' CAMS




Web-based Decision Support Tools Real-Time Decision Making and Long-term Planning









Regional Assessment of Cover Crop Residue Decomposition and N Release

Thapactal. (2022)

Regional Assessment of Cover Crop Residue Decomposition and N Release

Thapactal. (2022)

Regional Assessment of Cover Crop Residue Decomposition and N Release

Thapactal. (2022)

Regional Assessment of Cover Crop Residue Decomposition and N Release

Thapactal. (2022)

Regional Assessment of Cover Crop Residue Decomposition and N Release


Regional Assessment of Cover Crop Residue Decomposition and N Release

Regional Assessment of Cover Crop Residue Decomposition and N Release

Regional Assessment of Cover Crop Residue Decomposition and N Release

Regional Assessment of Cover Crop Residue Decomposition and N Release

Regional Assessment of Cover Crop Residue Decomposition and N Release



Precision Nitrogen Management with Cover crops CC-NCALC: (https://covercrop-ncalc.org) FEEDBA Home Location Soil Cover Crop Cash Crop Output Example: Legume ∨ Residue N Content Field name RESIDUE REMAINING (123 lb/ac) By $\begin{tabular}{ll} {\bf 4} & \begin{tabular}{ll} \begin{tabula$ (Clover, Crimson) • 63 lb/ac for surface residues. Holo-cellulose Cover crop N released and Corn N uptake over time **Dry Biomass** Cover Crop N Released (lb/ac)
Corn N uptake (lb/ac)
00
01 Cash crop recommended N rate after accounting for cover crop N credits. No Till (Surface)

Ongoing Tool Calibration

Cover crop treatments:


• cereal rye, legume, rye/legume, fallow


Nitrogen rates


• 0, 45, 90, 179, 269, and 358 lb ac⁻¹





19

















Crop, Weed, and Cover Crop Performance Mapping Bluekier Bluekier 27

Crop, Weed, and Cover Crop Performance Mapping

RGB frame
Species Semantic Segmentation
Species and Biomass Estimates

Depth frame



High Resolution, Open-access Image Repositories

BenchBot V1

Contributing sites

Contributing sites

Enterprising BenchBot V2 with Farm-ng







31

Image Repositories that are Portable to All Camera Sensors









٠.
























National Outreach Team

GROW Website

www.growiwm.org

About 475 subscribers:

- Farmers
- Crop Advisors
- Media Outlets
- Ag Scientists
- Government organizations
- ✓ Weed Management Toolbox
- ✓ Weed Identification Pages
- V IWM Definitions, Resources & Tools
- News Page updated weekly with IWM research & news
- Newsletter: The Weekly Update from GROW





Social Media

•Twitter: @GetRidOfWeeds

•YouTube: GROW IWM

• 45 videos posted in 2023

•Facebook: GROW IWM

•Instagram: @grow.iwm



